


Massive Transfusion Protocol



Children <30 kg:

Pack 1	Pack 2	Pack 3	Pack 4
2 units PRBCs	2 units PRBCs	2 units PRBCs	2 units PRBCs
2 unit	2 unit	2 unit	2 unit
1/4 SDP	1/4 SDP	1/4 SDP	1/4 SDP
	2 units cryoprecipitate	2 units cryoprecipitate	2 units cryoprecipitate

Children > 30 KG

Pack 1	Pack 2	Pack 3	Pack 4
4 units PRBCs	4 units PRBCs	4 units PRBCs	4 units PRBCs
4 unit FFP	4 unit FFP	4 unit FFP	4 unit FFP
1 SDP	1 SDP	1 SDP	1 SDP
	3 pack pre-pooled cryoprecipitate	3 pack pre-pooled cryoprecipitate	3 pack pre-pooled cryoprecipitate

Adjuncts: Factor VII a- initial dose of 30 micrograms/kg up to 1 mg

PEDIATRIC TRAUMA SOCIETY CLINICAL PRACTICE GUIDELINES DISCLAIMER STATEMENT

These guidelines have been supplied by a hospital as an example of a clinical practice guideline to provide clinicians at that institution with an analytical framework for the evaluation and treatment of a particular diagnosis or condition. These guidelines are not intended to establish a protocol for all patients with a particular condition, may not be replicable at other institutions, and are not intended to replace a clinician's clinical judgment. A clinician's adherence to these guidelines is voluntary. It is understood that some patients will not fit the clinical conditions contemplated by these guidelines and that the recommendations contained in these guidelines should not be considered inclusive of all proper methods or exclusive of other methods of care reasonably directed to obtaining the same results. Decisions to adopt any specific recommendation of these guidelines must be made by the clinician in light of available resources and the individual circumstances presented by the patient. These guidelines have not undergone expert review by the Pediatric Trauma Society and their posting by the Pediatric

Trauma Society should not be considered an endorsement of their content or the refutation of any alternate management strategy.